



29 Main Street, Truro, NS B2N 4G5

Phone: (902) 893-4331

Fax 897-4442

www.centralnovavet.com

FELINE CARDIOMYOPATHY

What is cardiomyopathy?

Literally, the term "cardiomyopathy" means disease of the heart muscle. More specifically, cardiomyopathy (CM) is a disease of the heart muscle in which either the heart walls thicken greatly (hypertrophic and restrictive forms) or stretch greatly (dilated form). In either form, the heart's function is significantly compromised and leads to an eventual state of heart failure.

What causes cardiomyopathy?

Several causes of feline cardiomyopathy have been identified. A deficiency of taurine, an essential amino acid, will cause dilated CM in the cat. This was a common occurrence many years ago, but now commercial cat foods are now properly supplemented. Hyperthyroidism, a non-cancerous growth of the thyroid gland, can cause a variation of the hypertrophic form of CM. The restrictive form is associated with an unidentified inflammatory process within the heart muscle. However, most cases of CM are not caused by any of these processes, and we do not understand their origin.



What does a cat with cardiomyopathy look like?

Cardiomyopathy is a disease that usually takes several weeks to months to progress to a serious stage. During the early weeks of the disease, the cat will probably look normal. Cats have the ability to hide serious illness until it reaches a crisis stage. Therefore, most cats that develop clinical signs of cardiomyopathy **will appear to have been ill for only a few days**. A few days of inactivity and poor appetite occur first. Just prior to the state of heart failure and death, the cat may become very inactive and exhibit **laboured breathing**. Both may be due to insufficient oxygen transport to the body's tissues; the latter may also be due to a collection of fluid in or around the lungs.

How is this disease diagnosed?

Diagnosis is generally made with a **chest radiograph (X-ray)**. The heart will have an abnormal shape and fluid in or around the lungs may be detected. If a large amount of fluid is present around the lungs, it may be necessary to remove it and take more X-rays because the presence of this fluid interferes with evaluation of the heart. Many cases also require better visualization of the heart with an **echocardiogram, (cardiac ultrasound)**. This is a non-

invasive method for looking at the heart while it is pumping. Referral to Halifax or the Atlantic Veterinary College is required for this. The x-ray can tell us about the size and shape of the heart but nothing about heart function. The ultrasound can provide information on function. The ultrasound will also allow measurement of the heart muscle to determine if it is too thick (hypertrophic CM) or too thin (dilated CM). Finally, an electrocardiogram (EKG) is useful to evaluate the rhythm of the heart.


Determination of the level of thyroid hormone in the blood (T_4) is often indicated in evaluating cats with hypertrophic CM. This simple blood test can help identify an overactive thyroid gland as the underlying cause of heart disease.

What is involved with treatment?

Treatment is based on the type of CM present & involved giving your pet oral medication (pills) to help the heart function, reduce fluid accumulation & prevent clot formation.. Unfortunately, we often diagnose this disease in its later stages, so some cats do not respond well to treatment. If treatment is successful, continual medication will be necessary since the disease cannot be cured. Cats with CM tend to have a shortened life-expectancy.

If hyperthyroidism is identified in a cat with hypertrophic CM, the heart disease is potentially reversible if the cat receives appropriate and timely treatment for the thyroid disease.

Hypertrophic Cardiomyopathy



Are there complications that may occur?

Most of the cats with cardiomyopathy develop signs of heart failure, as previously described. Also, cats with CM are prone to producing **blood clots** within their hearts. When these clots escape the heart, they travel through various arteries leading from the heart. They eventually lodge in a narrow part of the artery when their continued travel becomes impaired by the artery's diameter. The most common site for clots to lodge is the point at which the aorta splits before going into the rear legs. Thus, these cats often become **paralyzed** in the rear legs very suddenly and are in significant pain. In many cases, it is paralysis and pain that first becomes noticeable and is the reason that medical treatment is sought. Some owners mistake this event for an uncomplicated lameness, or even a broken leg. When these cats are examined, there are no pulses to one or both rear legs, the legs are cold, and the footpads appear blue (cyanotic) due to the lack oxygen.

Treatment of the paralyzed cat concentrates on drugs to relieve pain and to hasten the return of circulation to the legs. Since these cats also have severe heart disease the prognosis is variable, but is improved if the cat receives immediate attention. Within 3-10 days, circulation may improve. However, the heart disease must be controlled quickly or they will die.

What is the prognosis for cats with cardiomyopathy?

The prognosis for CM is quite variable, depending on the form of the disease and the severity at the time of diagnosis. Some cats can live for years if properly medicated, but the survival rate averages about six months. Again, the exception is when CM is caused by hyperthyroidism. If hyperthyroidism is successfully treated, the heart function may return to normal and the cat will no longer require treatment for heart disease.